

Allium ekimianum: a new species (Amaryllidaceae) from Turkey

Gülnur Ekşi¹, Mehmet Koyuncu², Ayşe Mine Gençler Özkan¹

1 Department of Pharmaceutical Botany, Faculty of Pharmacy, Ankara University, Ankara, Turkey **2** Faculty of Pharmacy, Cyprus International University, Haspolat-Lefkoşa, Cyprus

Corresponding author: Gülnur Ekşi (gulnur_eksi@yahoo.com)

Academic editor: C. Morden | Received 15 January 2016 | Accepted 28 March 2016 | Published 6 April 2016

Citation: Ekşi G, Koyuncu M, Özkan AMG (2016) *Allium ekimianum*: a new species (Amaryllidaceae) from Turkey. PhytoKeys 62: 83–93. doi: 10.3897/phytokeys.62.7796

Abstract

Allium ekimianum is described here as a new species. This taxon belongs to the genus *Allium* section *Allium* and grows in Elazığ Province (East Anatolia, Turkey). It is a narrowly distributed species and morphologically most similar to *A. asperiflorum* and *A. sintenisii*, and *A. erzincanicum* but it is clearly differentiated due to the curved stem, smooth pedicel surfaces, bracteole arrangements at pedicel bases, tepal lengths and surfaces. In this study, a comprehensive description, distribution map of *A. ekimianum*, identification key, and detailed illustrations are provided for *A. ekimianum* and related taxa.

Keywords

Allium, section *Allium*, endemic species, taxonomy, Turkey

Introduction

The genus *Allium* L. is one of the largest monocotyledonous genera with c. 900 species distributed world-wide (Govaerts et al. 2013, Keusgen et al. 2011). The genus was formerly included in the Liliaceae family, but the Angiosperm Phylogeny Group (APG) reassessed the taxonomic position of this genus and finally *Allium* was placed in the Amaryllidaceae family (APG III 2009). The primary evolution center of the genus extends across the Irano–Turanian biogeographical region, and the Mediterranean basin and western North America are secondary centres of diversity (Friesen et al. 2006).

Based upon these centres, *Allium* species have scattered widely all over the northern hemisphere (Hanelt 1990, Fritsch and Friesen 2002). The genus is characterized by having bulbs enclosed in membranous (sometimes finally fibrous) tunics, terminal umbel, free or almost free, 1-veined tepals, often a subgynobasic style and loculicidal capsule with one or two seeds per loculus (Kollmann 1984).

Following the results of recent molecular investigations, *Allium* is divided into 15 subgenera and 56 sections (Friesen et al. 2006). Subgenus *Allium* is the largest, comprising approximately 280 species (Hanelt et al. 1992), 114 of which compose its largest section, *Allium* (Mathew, 1996). Section *Allium* encompasses those species of *Allium* that have a well-developed bulb, stem (never basal) leaves, campanulate to cup-shaped (never stellate) flowers, and filaments in two distinct whorls, the outer three nearly always simple and the inner three markedly tricuspidate (rarely 5-7 cuspidate) with the anther attached to the median cusp. This section includes economically important species, such as garlic (*A. sativum* L.) and leek (*A. ampeloprasum* L.), as well as other minor crops of local importance, such as great headed garlic, and kurrat (Block 2010). Despite the major importance of the section *Allium*, it has not been subjected to a comprehensive molecular taxonomic evaluation; only partial molecular genetic studies that involved a limited set of species have been published (Kik et al. 1997; Havey and Leite 1999; Bohanec et al. 2005; Hirscheegger et al. 2010). Interspecific and infraspecific relationships within this section still remain unresolved. As reviewed by Mathew (1996), polyploidy is a common feature in section *Allium*. However, to a certain extent, it has been left unexplored, leaving the origin of polyploid species undetermined (Hirscheegger et al. 2010).

Turkey has approximately 190 *Allium* taxa in 14 sections, c. one-third endemic, demonstrating that it is a prominent part of the southeastern Asian center of *Allium* diversity (Ekşi et al. 2015; Özhatay and Kandemir 2015; Koyuncu 2012). Section *Allium* remains the most species-rich section of the genus (Friesen et al. 2006).

Turkey has four reasons for having an exceptionally rich flora. First, it is the meeting point of three phytogeographical regions, the Euro–Siberian, Mediterranean and Irano–Turanian regions. Second, Anatolia (Asian part of Turkey) is a passageway and a migration route between Southern Europe and the flora of South–West Asia allowing the penetration of Asiatic elements into South Europe. Third, many taxa have their center of origin and/or center of diversity in Anatolia. Fourth, the endemism ratio is high, presumably connected with the climatic and topographical diversity of the country (Davis 1965, 1971).

During the revision of the treatment of *Allium* in Turkey, individuals of a new species were collected by Prof. Dr. Mehmet Koyuncu in 1983 from Eastern Anatolia. They belong to *Allium* section *Allium* due to ovoid bulb, linear leaves, campanulate to ovoid perigon; 3-cuspidate inner filaments, distinct nectariferous pores on ovary, ovule numbers in per loculus (Kollmann 1984). The initial evaluation suggested this collection was a form of *A. asperiflorum* Misch. However, detailed examination of herbarium material and a review of the literature indicated this represented an undescribed species. The present study is focused on the morphological characters for distinguishing

a new species in *Allium* section *Allium*. Investigations on living and herbarium specimens suggest that this new species is morphologically most similar to *A. asperiflorum*, *A. erzincanicum* N. Özhatay & Kandemir and *A. sintenisii* Freyn.

Materials and methods

The overall morphology of the new species was examined by stereo binocular microscope (Leica Zoom 2000). For morphological comparisons, we consulted dry herbarium material kept in AEF, ANK, E, GAZI, and ISTE (acronyms according to Thiers 2015). The *Flora of Turkey* and floras of the neighbouring regions including Iran, Iraq, and Syria were also consulted (Boissier 1882, Feinbrun 1948, Wendelbo 1971, 1985, Kollmann et al. 1983, Kollmann 1984, Mathew 1996, Özhatay and Tzanoudakis 2000). The diagnostic traits of the new species and its three most similar relatives are presented in Table 1. Distribution map of compared taxa is provided according to the Grid classification system developed by Davis (1965) in Figure 1. IUCN conservation assessment and justification is proposed according to the criteria established by IUCN (IUCN 2003). In addition, an identification key is provided to distinguish new species and closely related taxa.

Taxonomic treatment

Allium ekimianum Ekşi, Koyuncu & Özkan, sp. nov.

urn:lsid:ipni.org:names:60471703-2

Figure 2

Note. Diagnostic characters for *Allium ekimianum* include curved stem, smooth pedicels, united bracteoles, verrucose–scabrid and straight outer tepal, smooth inner tepal, longer inner tepal.

Type. Turkey. Elazığ: Fırat University, steppe, c. 1150 m, 02.07.1983, *Koyuncu 7847* (holotype: AEF!, isotype: GAZI!).

Description. Bulb ovoid, 0.7–1.2 × 1–1.5 cm; outer tunics membranous, brownish, ± breaking into parallel fibres; inner tunics white; bulblets absent. Stem 15–35 cm, curved, often purplish below. Leaves 2–3, linear, 1–2 mm broad, flat, shorter than scape, sheathing lower ½ of stem. Umbel globose–subglobose, 1.5–3 cm diameter, dense, 20–60 flowered. Spathe caducous. Pedicels smooth, unequal, not elongating in fruit; up to 2.5 × perigon; bracteoles present, united at the base of outer pedicels, splitting into several lobes at apex, c. 5 mm. Perigon ovoid, campanulate; tepals purple, pale pink; outer tepals straight, 5 × c. 3.5 mm, obovoid, verrucose–scabrid, acute–subacute, obtus at apex; inner tepals c. 4 × 2 mm, narrowly oblong, smooth, obtus at apex. Stamens included; filaments ciliate at base; inner filaments 4 × 2 mm; median cusps c. 1 mm, slightly shorter than lateral cusps (c. 1.5 mm); basal lamina c. 3 mm, 3

Table 1. Main differences between *A. asperiflorum*, *A. ekimianum* and *A. sintenisii*, *A. erzincanicum*.

	<i>A. asperiflorum</i>	<i>A. ekimianum</i>	<i>A. sintenisii</i>	<i>A. erzincanicum</i>
Outer bulb tunics	membranous	membranous	membranous	reticulate-fibrous
Stem	erect	curved	almost erect	erect
Leaves	densely scabrid	glabrous	densely scabrid	glabrous or almost glabrous
Bracteoles	solitary at the base of each pedicel	outer ones with united bracteoles at base	outer ones with united bracteoles at base	outer ones with united bracteoles at base
Pedicel surface	papillose	smooth	scabrid near the base of perianth	smooth
Outer tepal	densely papillose, keeled	verrucose–scabrid, straight	loosely bearded with long white papillae at whole surface, keeled	loosely bearded with long white papillae just on the midvein, keeled
Inner tepal	papillose, 5–7 × 2.5 mm, equal to outer tepal	smooth, c. 4 × 2 mm, distinctly longer than outer tepal	smooth, 7–10 × 3 mm, almost equal to outer tepal	loosely bearded with long white papillae just on the midvein, 4–5 × 1–2 mm, equal to outer tepal




Figure 1. Distribution map of *A. ekimianum*, *A. asperiflorum*, *A. sintenisii*, *A. erzincanicum*.

times longer than median cusps. Anther 1 mm, yellow. Pistil c. 3–5 mm; style c. 1–2 mm; ovary c. 2–3 × 1.5–2 mm, ovoid, smooth. Capsule 4 × 3.5 mm, ovoid; valves emarginate–bilobate at apex; seed 3 × 1.5 mm, black.

Etymology. The species is named in honor of the eminent Turkish botanist Prof. Dr. Tuna Ekim, who dedicated his life to Turkish Flora, was retired from İstanbul University.

Distribution and ecology. The distribution of *Allium ekimianum* is restricted to Province of Elazığ from East Anatolia, where it grows on steppe between 1100–




Figure 2. *Allium ekimianum* (Koyuncu 7847/Elazığ). Plant (**A**, **B**), flower (**C**), flower longitudinal section (**D**), inner tepal (**D1**, **D2**), outer tepal (**D3**, **D4**), pistil (**E**), capsule (**F**), seed (**G**), leaf sheathing (**H**), leaf cross section (**I**). (Drawn by Gülnur Ekşi).

1200 m of elevation. Species associated with *A. ekimianum* include *Campanula stricta* L., *Silene italica* (L.) Pers., *Silene vulgaris* (Moench) Garcker, *Euphorbia macroclada* Boiss., *Papaver rhoeas* L., *Crataegus monogyna* Jaq., *Rosa canina* L., *Rosa × dumalis* Bechst., *Potentilla erecta* L., *Sanguisorba minor* Scop., *Achillea millefolium* L., *Allium*




Figure 3. *Allium asperiflorum* (Koyuncu 10539/Artvin). Plant (**A1, A2, B**), flower (**C**), flower longitudinal section (**D**), inner tepal (**D1, D2**), outer tepal (**D3, D4**), pistil (**E**), capsule (**F**), seed (**G**), bulblet (**H**), leaf surface (**I**), leaf sheathing (**J**). (Drawn by Gülnur Ekşi).




Figure 4. *Allium sintenisii* (Koyuncu 9692/Kahramanmaraş). Plant (**A1, A2**), flower (**B**), flower longitudinal section (**C**), inner tepal (**C1, C2**), outer tepal (**C3, C4**), pistil (**D**), capsule (**E**), seed (**F**), bulblet (**G**), leaf sheathing (**H**), leaf surface (**I**). (Drawn by Gülnur Ekşi).

scorodoprasum L., *Vicia cracca* L., *Crepis foetida* L., *Eryngium campestre* L., *Salvia verticillata* L., *Avena sterilis* L. Elazığ is located on the east of Anatolian diagonal, in the skirts of South-Eastern Taurus Mountains (Çakılcıoğlu et al. 2008), in the Upper Euphrates Region of the Eastern Anatolia Region (Şengün 2007). Elazığ belongs to the Irano-Turanian Plant Geography Region and falls within the B7 grid square (Davis 1965). The Irano-Turanian Region is confined to Central and East Anatolia. This great region of steppe, mountain steppe and semi-desert is also characterized by the existence of a hypothetical oblique line that runs from Bayburt-Gümüşhane southwardly to Anti-Taurus where it bifurcates with one prong leading to the Amanus and the other to the Cilician Taurus. This line is called “Anatolian Diagonal” (Figure 1). The flora of central Anatolia as the western side of the Diagonal is floristically different from the rest of the Irano-Turanian region to the east. According to the plant distribution patterns in eastern Anatolia, many endemics are restricted to part of the Diagonal belt, or extend right along it (Davis 1965, Davis 1971).

IUCN Conservation Assessment and Justification. Following the criteria established by IUCN (IUCN 2003), an initial provisional assessment of Critically Endangered (CR) (criteria B2a + B2biii) is suggested for this new taxon. This species occurs only in Elazığ University campus area in Elazığ province (East Anatolia) at 1100–1200 m. The area is under subversive people activities such as new constructions of buildings. As a result, the habitat of *Allium ekimianum* is highly threatened of vanishing by people activities. The area of *A. ekimianum* occupancy (AOO) is less than 10 km² with the number of mature individuals which is under reduction and being less than 50.

Related species. *Allium ekimianum* is closely related to *A. asperiflorum* and *A. sintenisii* and *A. erzincanicum*. All four species share traits of ovoid bulb, globose to subglobose umbel, campanulate to ovoid perigon, rough outer tepal surfaces, stamens shorter than perigon, ovoid ovary. *Allium ekimianum* differs from *A. asperiflorum*, *A. sintenisii* and *A. erzincanicum* in its outer tunics, stem, leaves, bracteoles, pedicel surface, outer tepal, and inner tepal characters. The three species are compared in Table 1 and these traits are illustrated in Figures 2–4.

Key for identification of *Allium* species related to *A. ekimianum*

- | | | |
|---|--|------------------------|
| 1 | Outer perianth segments loosely bearded at whole surface or along midvein with long white papillae, pale pink, more intensely pink at tip..... | 2 |
| – | Outer perianth segments not bearded on whole surface or along midvein... | 3 |
| 2 | Perigon 7–10 mm; outer tunics membranous; leaves scabrid; pedicels scabrid near the base of perianth; outer tepal loosely bearded with long white papillae at whole surface; inner tepal smooth | <i>A. sintenisii</i> |
| – | Perigon 4–5 mm; outer tunics reticulate fibrose; leaves glabrous or almost glabrous; pedicels smooth; outer tepal loosely bearded with long white papillae just on the midvein; inner tepal bearded on the midvein | <i>A. erzincanicum</i> |

- 3 Stem erect; involucre-like structure absent; outer tepal keeled, surface densely papillose; inner tepal surface scarcely papillose; leaves densely scabrid; pedicels densely papillose; bulblets numerous ***A. asperiflorum***
- Stem curved; involucre-like structure present; outer tepal not keeled, surface verrucose-scabrid; inner tepal surface glabrous; leaves glabrous; pedicels smooth; bulblets ± present ***A. ekimianum***

Additional specimens examined

The capital letters and the numbers in bold after species names represent the Grid classification system (Davis, 1965) and the names in bold are the provinces from the Eastern Turkey. The abbreviations and the numbers in brackets at the end of the sentences represent the herbarium names and the accession numbers, respectively.

Allium asperiflorum: **A9 Artvin**: Borçka–Artvin arası, Artvin’e 10 km kala, kayalıklar, 200 m, 14 vii 1993, *M. Koyuncu* 10539 (AEF 18113). **B6 Sivas**: Divriği–Cürekarası, 3 vi 1983, *H. Başer s.n.* (ESSE 3320). **B7 Erzincan**: İliç, Hassanova village, 900–1100 m, *Çelik s.n.* (AEF 5699). İliç, Hasanova Köyü altındaki Tepeler, 900–1000, 16 vi 1976, *N. Çelik s.n.* (AEF 5699). Erzincan–Refahiye yolu, 35 km, kuruçakıllı yamaçlar, 1900 m, 22 viii 1990, *M. Koyuncu* 8808 (AEF 15737). İliç–Refahiye çevresi, Gümüşahar’dan sonar Sunibeli Geçiti, orman açıklıkları, 1700 m, 21 vi 2005, *M. Koyuncu* 15098 & *N. Aslan* (AEF 24263). **B7 Tunceli**: Ovacık üzeri, Munzur Dağı, Kepir Gediği, kayalık taşlık arazi, 2400–2750 m, 10 viii 1976, *M. Koyuncu* & *N. Çelik s.n.* (AEF 5683). **C8 Siirt**: Pervari’nin üstü, kalker kayalıklar, stepler, 1600–1700 m, 16 vi 1980, *M. Koyuncu* 3260 (AEF 9563). ***Allium sintenisii***: **B6 Kayseri**: Sarız, Yalak, Binboğa Dağı, 2000–2200 m, 1 vii 1992, *M. Koyuncu* & *H. Duman* 5175 (AEF 17830). Bakır Dağı–Tufanbeyli arası, Gezbeli Geçidi, 2200 m, 28 vii 2008, *M. Koyuncu* 15993 (AEF 25277). **B6 Malatya**: Kuluncak, Kızılyüce Dağı kuzey eteği, çayırılık, 1700 m, 18 vi 1994, *B. Yıldız* 11582 (AEF 26253). **B6 Maraş**: Göksun–Binboğa Dağı, 2000–2400 m, 17 vii 1992, *M. Koyuncu* 9692, *H. Duman*, *Z. Aytaç* (AEF 17465). **B7 Malatya/Sivas**: Kangal to Hekimhan, 1300 m, *Stn.* & *Hend.* 5390. **B7 Erzincan**: Spikör Dağı Geçidi, step yamaçlar, 2300 m, 27 viii 2008, *M. Koyuncu* 16128 (AEF 25453) **ibid.** 14 vii 2009, *M. Koyuncu* 16176 (AEF 25593). ***Allium erzincanicum***: **B7 Erzincan**: Munzur Dağları, Mercan Suyu, kalker kayalıklar, 37 S 550540 D, 4374863 K, 1997 m, 08 vii 2014, *Kandemir* 10613 (isotype NGBB).

Acknowledgement

We would like to thank Dr. Sabina Knees for help in improving to manuscript. This study is financially supported by the Edinburgh Botanic Garden (SIBBALD) Trust and the Scientific and Technological Research Council of Turkey (TÜBİTAK). We also would like to thank to AEF, ANK, E, GAZI, ISTE and NGBB herbaria for their kind support.

References

- APG (2009) An update of the angiosperm phylogeny group classification for the orders and families of flowering plants: APG III. *Botanical Journal of Linnean Society* 161: 105–121. doi: 10.1111/j.1095-8339.2009.00996.x
- Block (2010) *Garlic and Other Alliums – The Lore and The Science*. The Royal Society of Chemistry, Cambridge, 480 pp.
- Bohanec M, Messean A, Scatasta S, Dzeroski S, Znidarsic M (2005) A qualitative multi-attribute model for economic and ecological evaluation of genetically modified crops. *EnviroInfo Brno 2005: Proceedings of the 19th International Conference on Informatics for Environmental Protection*, Masaryk University in Brno, 661–668.
- Boissier E (1882) *Flora Orientalis* Volume 5. Geneve & Basel.
- Çakılçioğlu U, Türkoğlu I, Kürşat M (2008) The flora of Çitli Lowland (Elazığ). *e-Journal of New World Sciences Academy* 3: 232–249.
- Davis PH (1965) *Flora of Turkey and the East Aegean Islands* (Volume 1). Edinburgh University Press, Edinburgh, 568 pp.
- Davis PH (1971) Distribution patterns in Anatolia with particular reference to endemism. In: Davis PH, Harper PC, Hedge IC (Eds) *Plant Life of South–West Asia*. The Botanical Society of Edinburgh, Edinburgh, 15–27.
- Ekşi G, Koyuncu M, Bona M (2015) *Allium phaneranthum* subsp. *involucratum* (Amaryllidaceae), a new species from Turkey. *Bangladesh Journal of Plant Taxonomy* 22(2): 143–146. doi: 10.3329/bjpt.v22i2.26076
- Feinbrun N (1948) Further studies on *Allium* of Palestine and the neighbouring countries. *Palestine Journal of Botany* 4: 144–157.
- Friesen N, Fritsch RM, Blattner FR (2006) Phylogeny and new intra generic classification of *Allium* (Alliaceae) based on nuclear ribosomal DNA ITS sequences. *Aliso* 22: 372–395.
- Fritsch RM, Friesen N (2002) Evolution, domestication and taxonomy. In: Rabinowitch HD, Currah L (Eds) *Allium crop science: recent advances*. CABI publishing, Oxfordshire, 5–30. doi: 10.1079/9780851995106.0005
- Hanelt P (1990) Taxonomy, evolution, and history. In: Rabinowitch HD, Brewster JL (Eds) *Onions and allied crops* (Volume 1). CRC Press, Boca Raton, 1–26.
- Hanelt P, Schultze-Motel J, Fritsch R, Kruse J, Maass HI, Ohle H, Pistrick K (1992) Infrageneric Grouping of *Allium*. The Gatersleben Approach. In: Hanelt P, Hammer K, Knupffer (Eds) *The genus Allium. Taxonomic Problems and Genetic Resources*, 107–123.
- Havey MJ, Leite DL (1999) Toward the Identification of Cytoplasmic Male Sterility in Leek: Evaluation of Organellar DNA Diversity among Cultivated Accessions of *Allium ampeloprasum*. *Journal of the American Society for Horticultural Science* 124(2): 163–165.
- Hirschegger P, Jakše J, Trontelj P, Bohanec B (2010) Origins of *Allium ampeloprasum* horticultural groups and a molecular phylogeny of the section *Allium* (*Allium*: Alliaceae) *Molecular Phylogenetics and Evolution* 54: 488–497. doi: 10.1016/j.ympev.2009.08.030
- Govaerts R, Kington S, Friesen N, Fritsch R, Snijman DA, Marcucci R, Silverstone-Sopkin PA, Brullo S (2013) World checklist of Amaryllidaceae. <http://apps.kew.org/wcsp/>

- IUCN (2003) Guidelines for Application of IUCN Red List Criteria at Regional Levels (Version 3). IUCN, Gland and Cambridge.
- Keusgen M, Kusterer J, Fritsch RM (2011) *Allium* species from Middle and Southwest Asia are a rich source for Marasmin. *Journal of Agricultural and Food Chemistry* 59: 8289–8297. doi: 10.1021/jf201052u
- Kik C, Samaylov AM, Verbeek WHJ, van Raamsdonk LWD (1997) Mitochondrial DNA variation and crossability of leek (*Allium porrum*) and its wild relatives from the *Allium ampeloprasum* complex. *Theoretical and Applied Genetics* 96: 465–471. doi: 10.1007/s001220050438
- Kollmann F, Özhatay N, Koyuncu M (1983) New *Allium* taxa From Turkey. *Notes From The Royal Botanic Garden Edinburgh* 41: 263.
- Kollmann F (1984) *Allium* L. In: Davis PH (Ed.) *Flora of Turkey and the East Aegean Islands*. Volume 8. Edinburgh University Press, Edinburgh, 98–211.
- Koyuncu M (2012) *Allium* L. In: Güner A, Aslan S, Ekim T, Vural M, Babaç MT (Eds) *Türkiye Bitkileri Listesi (Damarlı Bitkiler)*. Nezahat Gökyiğit Botanik Bahçesi ve Flora Araştırmaları Derneği Yayını, İstanbul, 30–44.
- Mathew B (1996) Review of *Allium* Section *Allium*. Royal Botanic Gardens, Kew, 182 pp.
- Özhatay N, Kandemir A (2015) *Allium erzincanicum* (sect. *Allium*): Doğu Anadolu Bölgesi'nden yeni bir *Soğan* (*Allium* L.) türü. *Bağbahçe Bilim Dergisi* 1(2): 50–58.
- Özhatay N, Tzanoudakis D (2000) *Allium* L. In: Güner A, Özhatay N, Ekim T, Başer KHC (Eds) *Flora of Turkey and the East Aegean Islands Volume 11*. Edinburgh University Press, Edinburgh, 224–232.
- Thiers B (2015) Index Herbariorum: A global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. Available from: <http://sweetgum.nybg.org/ih/> [accessed 4 September 2015]
- Wendelbo P (1971) *Allium* L. In: Rechinger KH (Ed.) *Flora Iranica Volume 76*. Akademische Druck und Verlagsanstalt, Graz, 1–100.
- Wendelbo P (1985) *Allium* L. In: Townsend CC, Guest E (Eds) *Flora of Iraq Volume 8*. Ministry of Agriculture Republic of Iraq, Baghdad, 137–177.